

Greenberger-Horne-Zeilinger nonlocality for continuous variable systems

Zeng-Bing Chen¹ and Yong-De Zhang^{2,1}

¹*Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230027, P.R. China*

²*CCAST (World Laboratory), P.O. Box 8730, Beijing 100080, P.R. China*

(February 1, 2008)

As a development of our previous work, this paper is concerned with the Greenberger-Horne-Zeilinger (GHZ) nonlocality for continuous variable cases. The discussion is based on the introduction of a pseudospin operator, which has the same algebra as the Pauli operator, for each of the N modes of a light field. Then the Bell-CHSH (Clauser, Horne, Shimony and Holt) inequality is presented for the N modes, each of which has a continuous degree of freedom. Following Mermin's argument, it is demonstrated that for N -mode parity-entangled GHZ states (in an infinite-dimensional Hilbert space) of the light field, the contradictions between quantum mechanics and local realism grow exponentially with N , similarly to the usual N -spin cases.

PACS number(s): 03.65.Ud, 03.65.Ta

Under the presumption of local realism, Einstein, Podolsky and Rosen (EPR) argued that it seems possible to assign values with certainty to canonically conjugate variables (e.g., position and momentum) of one particle by performing experiments on another particle of an entangled biparticle system [1]. EPR's observation is definitely in conflict with quantum mechanics. The entangled states of the biparticle system are now known as the EPR states of continuous variables. In 1965 Bell, using Bohm's EPR states of discrete variables (e.g., spin) [2], derived the famous Bell's inequalities [3–5], enabling quantitative tests of quantum mechanics versus local realism. So far, many experiments based on Bohm's EPR states completely confirmed quantum mechanics [6]. Bell's theorem without inequalities has also been demonstrated for nonmaximally entangled biparticle states [7] and multiparticle Greenberger-Horne-Zeilinger (GHZ) states [8,9]. In this case, the contradictions between quantum mechanics and local realistic theories are of a non-statistical nature; the nonlocality can be manifest in a single measurement. Using three-photon polarization-entangled states, Pan *et al.* carried out an experimental test of GHZ theorem [10], confirming again the correctness of quantum mechanics. Besides its theoretical interest, research work on quantum nonlocality also gains great impetus from the rapidly developing field of quantum information theory [11].

Quantum nonlocality for continuous variables has also attracted much attention recently [5,12–14]. Several proposals have been put forward to prepare the original EPR states [15–17]; EPR paradox in EPR's original sense has been realized experimentally [18,19]. The experiment exploited the nondegenerate optical parametric amplifier (NOPA), which generates the two-mode squeezed vacuum states (“regularized” EPR states)

$$|\text{NOPA}\rangle = \frac{1}{\cosh r} \sum_{n=0}^{\infty} (\tanh r)^n |n, n\rangle, \quad (1)$$

with $r > 0$ denoting the squeezing parameter and $|nn\rangle \equiv |n\rangle_1 \otimes |n\rangle_2$ the twin-photon states. In the limit of infinite squeezing, the states $|\text{NOPA}\rangle$ reduce to a normalized version of the original EPR states [18,19]. Since the Wigner function [20] of $|\text{NOPA}\rangle$ is positive everywhere, it has been argued that the original EPR states may allow a hidden variable description and thus will not exhibit nonlocality [5,12,18,19]. This point of view was dramatically changed by Banaszek and Wódkiewicz (BW) [13]. Within a phase space formulation, BW used the “Bell operator” [21] based on the joint parity measurements and demonstrated the violations of the Bell inequality due to Clauser, Horne, Shimony and Holt (CHSH) [4]. GHZ nonlocality for continuous variables has also been analyzed [22], relying on the BW formalism. On the experimental aspect, a recent experiment [23] observed, for the first time, the violations of the Bell-CHSH inequality by the regularized EPR states produced in a pulsed NOPA within the homodyne detection scheme.

In a recent paper [14], we have generalized Bell's inequalities to the continuous variable cases for biparticle systems. Our formalism is based on a new Bell operator, which is a direct analogy of its discrete variable counterpart [21]. In such a physically appealing manner, Bell's inequalities for both discrete variable (in a two-dimensional Hilbert space) and continuous variable (in an infinite-dimensional Hilbert space) cases take a mathematically similar form. In this paper, we extend the work to entangled multiparty systems. In particular, we will present GHZ theorem for the parity-entangled GHZ states of a three-mode light field. Moreover, the Bell-CHSH inequality for multiparty systems of discrete variables [24] will be generalized to continuous variable cases.

Let us start with considering a light field with three independent modes 1, 2 and 3. For each mode of the light field, we can introduce the “parity spin” operator (i.e., pseudospin operator acting upon the parity space of photons) [14]

$$\begin{aligned}\hat{\mathbf{s}}_j &= (s_{jx}, s_{jy}, s_{jz}), \quad (j = 1, 2, 3) \\ s_{jx} &= s_{j+} + s_{j-}, \quad s_{jy} = -i(s_{j+} - s_{j-}).\end{aligned}\quad (2)$$

Here

$$\begin{aligned}s_{jz} &= \sum_{n=0}^{\infty} \left[|2n\rangle_{jj} \langle 2n| - |2n+1\rangle_{jj} \langle 2n+1| \right], \\ s_{j+} &= \sum_{n=0}^{\infty} |2n\rangle_{jj} \langle 2n+1| = (s_{j-})^\dagger\end{aligned}\quad (3)$$

with $|n\rangle_j$ denoting the usual Fock states of the j -mode light field. After a simple algebra, it can be shown that

$$\begin{aligned}[s_{iz}, s_{j\pm}] &= \pm 2\delta_{ij}s_{i\pm}, \quad [s_{i+}, s_{j-}] = \delta_{ij}s_{iz}, \\ s_{i\mu}s_{j\nu} + s_{i\nu}s_{j\mu} &= 2\delta_{ij}\delta_{\mu\nu}. \quad (\mu, \nu = x, y, z)\end{aligned}\quad (4)$$

The commutation relations in Eq. (4) are identical to their counterparts of the spin-1/2 systems. Thus each parity spin operator $\hat{\mathbf{s}}_j$ acts as the counterpart of the Pauli operator σ . In particular the following relation is valid

$$\begin{aligned}U_j(\zeta_j, \hat{\mathbf{n}}_j) &= \exp\left(-i\frac{\zeta_j}{2}\hat{\mathbf{n}}_j \cdot \hat{\mathbf{s}}_j\right) \\ &= \cos\frac{\zeta_j}{2} - i\hat{\mathbf{n}}_j \cdot \hat{\mathbf{s}}_j \sin\frac{\zeta_j}{2}\end{aligned}\quad (5)$$

for $\hat{\mathbf{s}}_j$ as for σ . Here $\hat{\mathbf{n}}_j$ is an arbitrary unit vector. $U_j(\zeta_j, \hat{\mathbf{n}}_j)$ are, in fact, the rotation operators in the parity space of photons.

Now one can introduce states of definite parity for each mode

$$|+\rangle_j \equiv \sum_{n=0}^{\infty} \mathcal{A}_n^{(j)} |2n\rangle_j, \quad |-\rangle_j \equiv \sum_{n=0}^{\infty} \mathcal{A}_n^{(j)} |2n+1\rangle_j, \quad (6)$$

where $\sum_{n=0}^{\infty} |\mathcal{A}_n^{(j)}|^2 = 1$. Some important properties of these parity states are

$$\begin{aligned}s_{jz} |\pm\rangle_j &= \pm |\pm\rangle_j, \\ s_{j\pm} |\pm\rangle_j &= 0, \quad s_{j\pm} |\mp\rangle_j = |\pm\rangle_j.\end{aligned}\quad (7)$$

Hence $|\pm\rangle_j$ are the eigenstates of the parity operators s_{jz} , with the corresponding eigenvalues ± 1 ; the operators $s_{j\pm}$ “flip” the parity of $|\mp\rangle_j$ and annihilate $|\pm\rangle_j$. In this aspect, $|\pm\rangle_j$ are similar to the spin-up and spin-down states of the spin-1/2 systems. Using the rotation operators as in Eq. (5), one can construct eigenstates of $\hat{\mathbf{n}}_j \cdot \hat{\mathbf{s}}_j$ for each mode. Now the parity-entangled GHZ state of the light field can be directly constructed as

$$|\text{GHZ}\rangle_3 = \frac{1}{\sqrt{2}} (|+\rangle_1 |+\rangle_2 |+\rangle_3 - |-\rangle_1 |-\rangle_2 |-\rangle_3). \quad (8)$$

Note that this state is defined in an infinite-dimensional Hilbert space.

Following GHZ’s argument [8,9,11], it is directly verifiable that the GHZ state (8) satisfies the eigenvalue equations as

$$\begin{aligned}s_{1x}s_{2x}s_{3x} |\text{GHZ}\rangle_3 &= -|\text{GHZ}\rangle_3, \\ s_{1x}s_{2y}s_{3y} |\text{GHZ}\rangle_3 &= |\text{GHZ}\rangle_3, \\ s_{1y}s_{2x}s_{3y} |\text{GHZ}\rangle_3 &= |\text{GHZ}\rangle_3, \\ s_{1y}s_{2y}s_{3x} |\text{GHZ}\rangle_3 &= |\text{GHZ}\rangle_3.\end{aligned}\quad (9)$$

Thus quantum mechanically, one can determine any operator $(s_{1x}, s_{1y}, s_{2x}, \dots)$ by performing appropriate measurements on the other two modes of the light field. To establish a local realistic interpretation of the quantum-mechanical results (9), one assumes that the individual value of any operator $(s_{1x}, s_{1y}, s_{2x}, \dots)$ is predetermined, e.g., by local hidden variables, regardless of any set of parity spin measurements on the three modes of the light field. These predetermined values are denoted by $p_{1x}, p_{1y}, p_{2x}, \dots$, with $p_{jx}, p_{jy} = \pm 1$. To be consistent with Eq. (9), local realistic theories impose the following relations

$$\begin{aligned}p_{1x}p_{2x}p_{3x} &= -1, \quad p_{1x}p_{2y}p_{3y} = +1, \\ p_{1y}p_{2x}p_{3y} &= +1, \quad p_{1y}p_{2y}p_{3x} = +1.\end{aligned}\quad (10)$$

But in fact, these relations are not mutually consistent. For example, the latter three relations in Eq. (10) give $p_{1x}p_{2x}p_{3x}p_{1y}^2p_{2y}^2p_{3y}^2 = p_{1x}p_{2x}p_{3x} = +1$, which conflicts with the first relation in Eq. (10). Here we have used the fact that $p_{jy}^2 = 1$. Thus quantum-mechanical predictions are incompatible with local realistic theories for the parity-entangled GHZ states (8). Also, the contradictions between quantum mechanics and local realism occur for perfect correlations (i.e., for non-statistical predictions) of the three-mode light field. It is worth pointing out that the GHZ paradox associated with position-momentum variables has also been considered in terms of the Weyl algebra in Ref. [25].

In a real experiment, the perfect correlation condition in the GHZ argument is very difficult to implement practically. To face this difficulty, one has to rely on the multiparty generalization of Bell’s inequalities, which has been derived originally for discrete variables [24]. Here we generalize the multiparty Bell-CHSH inequality to the present continuous variable case.

First, let us define the Bell operator in the case of a two-mode light field [14]:

$$\begin{aligned}\mathcal{B}_2 &= (\mathbf{a}_1 \cdot \hat{\mathbf{s}}_1) \otimes [(\mathbf{a}_2 \cdot \hat{\mathbf{s}}_2) + (\mathbf{a}'_2 \cdot \hat{\mathbf{s}}_2)] \\ &\quad + (\mathbf{a}'_1 \cdot \hat{\mathbf{s}}_1) \otimes [(\mathbf{a}_2 \cdot \hat{\mathbf{s}}_2) - (\mathbf{a}'_2 \cdot \hat{\mathbf{s}}_2)],\end{aligned}\quad (11)$$

where $\mathbf{a}_{1,2}$ and $\mathbf{a}'_{1,2}$ are four unit vectors. The commutation relations in Eq. (4) lead to, e.g.,

$$(\mathbf{a}_1 \cdot \hat{\mathbf{s}}_1)^2 = I_1, \quad (12)$$

where I_1 is the unit operator for the mode-1 of the light field. Equation (12) implies that the eigenvalues of the

Hermitian operator $\mathbf{a}_1 \cdot \hat{\mathbf{s}}_1$ are ± 1 . Now the N -mode Bell operator can be recursively defined by

$$\begin{aligned} \mathcal{B}_N &= \mathcal{B}_{N-1} \otimes \frac{1}{2}[(\mathbf{a}_N \cdot \hat{\mathbf{s}}_N) + (\mathbf{a}'_N \cdot \hat{\mathbf{s}}_N)] \\ &\quad + \mathcal{B}'_{N-1} \otimes \frac{1}{2}[(\mathbf{a}_N \cdot \hat{\mathbf{s}}_N) - (\mathbf{a}'_N \cdot \hat{\mathbf{s}}_N)], \end{aligned} \quad (13)$$

where $\hat{\mathbf{s}}_m$ ($m = 1, 2, \dots, N$) are the parity spin operators as defined in Eq. (2), $\mathbf{a}_m$ and $\mathbf{a}'_m$ are all unit vectors. $\mathcal{B}'_N$ defined in Eq. (13) has the same expression as $\mathcal{B}_N$, but with $\mathbf{a}_m \leftrightarrow \mathbf{a}'_m$. Similarly to the discrete variable case [24], local realistic theories must satisfy the following N -party Bell-CHSH inequality

$$|\langle \mathcal{B}_N \rangle| \leq 2, \quad (14)$$

where $\langle \mathcal{B}_N \rangle$ is the expectation value of $\mathcal{B}_N$ with respect to any N -mode state of the light field.

Meanwhile, an upper bound of $|\langle \mathcal{B}_N \rangle|$ can be determined quantum mechanically. By analogy to the N -party Bell-CHSH inequality for discrete variables [24], we can prove, using the properties of $\hat{\mathbf{s}}_m$ [see Eq. (4)], that

$$\begin{aligned} \mathcal{B}_N^2 &= \mathcal{B}_{N-1}^2 \otimes \frac{1}{2}(1 + \mathbf{a}_N \cdot \mathbf{a}'_N) \\ &\quad + \mathcal{B}_{N-1}'^2 \otimes \frac{1}{2}(1 - \mathbf{a}_N \cdot \mathbf{a}'_N) \\ &\quad + [\mathcal{B}_{N-1}, \mathcal{B}'_{N-1}] \otimes \frac{i}{2}(\mathbf{a}_N \times \mathbf{a}'_N) \cdot \hat{\mathbf{s}}_N \\ &\leq (1 + |\mathbf{a}_N \times \mathbf{a}'_N|) |\mathcal{B}_{N-1}^2| \leq 2^{N+1}. \end{aligned} \quad (15)$$

In the last step of Eq. (15), we have used the fact that $\mathcal{B}_2^2 \leq 2^{2+1}$ [14], from which the upper bound 2^{N+1} of $\mathcal{B}_N^2$ can be recursively obtained. Therefore quantum mechanics gives an upper bound $2^{(N+1)/2}$ of $|\langle \mathcal{B}_N \rangle|$; N -mode entangled states of the light field can violate the N -mode Bell-CHSH inequality (14) by a maximal factor of $2^{(N-1)/2}$, similarly to the discrete variable case [24]. For the two-mode Bell-CHSH inequality, the upper bound $2^{3/2}$ of $|\langle \mathcal{B}_N \rangle|$ is known as the Cirel'son bound [26]. Interestingly, the N -party Bell-CHSH inequality (14), though expressed in terms of discrete variable operators (i.e., the parity spin operators), can be exploited to uncover the GHZ nonlocality of continuous variable states, e.g., the multi-mode entangled states considered in Ref. [22].

Can an N -mode entangled state of the light field maximally violate the N -mode Bell-CHSH inequality (14)? To answer this question, we follow Mermin's reasoning [24] and take, as an example, the following N -mode parity-entangled GHZ state (instead of the multi-mode entangled states as in Ref. [22])

$$|\text{GHZ}\rangle_N = \frac{1}{\sqrt{2}}(|+\rangle_1 |+\rangle_2 \cdots |+\rangle_N - |-\rangle_1 |-\rangle_2 \cdots |-\rangle_N). \quad (16)$$

From Eq. (7) we have

$$\begin{aligned} (s_{mx} \pm i s_{my}) |\mp\rangle_m &= 2 |\pm\rangle_m, \\ (s_{mx} \pm i s_{my}) |\pm\rangle_m &= 0. \end{aligned} \quad (17)$$

Then it can be easily verified that $|\text{GHZ}\rangle_N$ is an eigenstate of the following operator

$$\begin{aligned} A &= \frac{1}{2} \left[\prod_{m=1}^N (s_{mx} + i s_{my}) + \prod_{m=1}^N (s_{mx} - i s_{my}) \right], \\ A |\text{GHZ}\rangle_N &= -2^{N-1} |\text{GHZ}\rangle_N \end{aligned} \quad (18)$$

with eigenvalue -2^{N-1} . Hence, quantum mechanics predicts that

$$|{}_N \langle \text{GHZ} | A | \text{GHZ} \rangle_N| = 2^{N-1}. \quad (19)$$

By contrast, local realistic theories predict a much lower bound on the measured results of A for large N . Under the notion of local realism, there are predetermined values $p_{1x}, p_{1y}, p_{2x}, \dots, (p_{jx}, p_{jy} = \pm 1)$ corresponding to operators $s_{1x}, s_{1y}, s_{2x}, \dots$. Each of these predetermined values of these operators does not depend on other measurements. Consequently, local realistic prediction on the upper bound of A reads

$$\left| \text{Re} \prod_{m=1}^N (p_{mx} + i p_{my}) \right| \leq \left| \prod_{m=1}^N (\pm 1 \pm i) \right| = 2^{N/2}. \quad (20)$$

Compared with the quantum-mechanical prediction (19), local realism predicts a bound of A lower by a large factor $2^{N/2-1}$, growing exponentially with N .

As comparison, within the BW formalism the contradictions between quantum mechanics and local realism also grow with increasing number of parties for multi-mode entangled states generated from squeezed states and beam splitters [22]. But the growth seems to decrease for larger number of parties, instead of the exponential growth. In this aspect, the merit of the present formalism is manifest.

To summarize, we have discussed the GHZ nonlocality for continuous variables, as a natural development of our previous work [14]. The introduction of the parity spin operators satisfying the same algebra as the Pauli operator is crucial to our discussion. Then the N -party Bell operator of continuous variables was constructed in terms of the parity spin operators of the N -mode light field, resulting in a generalization of the multiparty Bell-CHSH inequality to the continuous variable cases. Following Mermin's argument, it has been shown that for N -mode parity-entangled GHZ states (in an infinite-dimensional Hilbert space) of the light field, the contradictions between quantum mechanics and local realism grow exponentially with N , similarly to the usual N -spin cases. Within our formulation, there is a striking similarity between the multiparty Bell theorem for both discrete and continuous variables. Our result shows that only partial (i.e., parity spin) information of the whole Hilbert space of continuous variables is sufficient for the mere

purpose of uncovering quantum nonlocality. On an experimental aspect, however, the problem of realizing the parity-entangled GHZ states remains open.

Note added.—After the completion of the work, we became aware of a related paper by Massar and Pironio [27]. These authors considered the GHZ paradox for continuous variables in terms of modular and binary variables.

We thank Rob Clifton for bringing Ref. [25] into our attention. This work was supported by the National Natural Science Foundation of China under Grants No. 10104014, No. 19975043 and No. 10028406 and by the Chinese Academy of Sciences.

- [22] P. van Loock and S. L. Braunstein, Phys. Rev. A **63**, 022106 (2001); see also quant-ph/0006029.
- [23] A. Kuzmich, I. A. Walmsley, and L. Mandel, Phys. Rev. Lett. **85**, 1349 (2000).
- [24] D. N. Klyshko, Phys. Lett. A **172**, 399 (1993); N. D. Mermin, Phys. Rev. Lett. **65**, 1838 (1990); S. M. Roy and V. Singh, Phys. Rev. Lett. **67**, 2761 (1991); N. Gisin and H. Bechmann-Pasquinucci, Phys. Lett. A **246**, 1 (1998).
- [25] R. Clifton, Phys. Lett. A **271**, 1 (2000).
- [26] B. S. Cirel'son, Lett. Math. Phys. **4**, 93 (1980).
- [27] S. Massar and S. Pironio, Phys. Rev. A **64**, 062108 (2001); quant-ph/0103048.

-
- [1] A. Einstein, B. Podolsky and N. Rosen, Phys. Rev. **47**, 777 (1935).
 - [2] D. Bohm, *Quantum Theory* (Prentice Hall, Englewood Cliffs, NJ, 1951).
 - [3] J. S. Bell, Physics (Long Island) **1**, 195 (1964).
 - [4] J. F. Clauser, M. A. Horne, A. Shimony and R. A. Holt, Phys. Rev. Lett. **23**, 880 (1969).
 - [5] J. S. Bell, *Speakable and Unspeakable in Quantum Mechanics* (Cambridge Univ. Press, Cambridge, 1987).
 - [6] A. Aspect, Nature (London) **398**, 189 (1999).
 - [7] L. Hardy, Phys. Rev. Lett. **71**, 1665 (1993).
 - [8] D. M. Greenberger, M. A. Horne, and A. Zeilinger, in *Bell's Theorem, Quantum Theory, and Conceptions of the Universe*, edited by M. Kafatos (Kluwer Academic, Dordrecht, 1989).
 - [9] D. M. Greenberger, M. A. Horne, A. Shimony, and A. Zeilinger, *Am. J. Phys.* **58**, 1131 (1990).
 - [10] J. W. Pan, D. Bouwmeester, M. Daniell, H. Weinfurter, and A. Zeilinger, *Nature* (London) **403**, 515 (2000).
 - [11] D. Bouwmeester, A. Ekert, and A. Zeilinger (eds), *The Physics of Quantum Information* (Springer-Verlag, Berlin, 2000).
 - [12] J. S. Bell, Ann. (N.Y.) Acad. Sci. **480**, 263 (1986).
 - [13] K. Banaszek and K. Wódkiewicz, Phys. Rev. A **58**, 4345 (1998); Phys. Rev. Lett. **82**, 2009 (1999).
 - [14] Z. B. Chen, J. W. Pan, G. Hou, and Y. D. Zhang, Phys. Rev. Lett. **88**, 040406 (2002).
 - [15] M. D. Reid and P. D. Drummond, Phys. Rev. Lett. **60**, 2731 (1988); M. D. Reid, Phys. Rev. A **40**, 913 (1989).
 - [16] D. F. Walls and G. J. Milburn, *Quantum Optics* (Springer-Verlag, Berlin, 1994).
 - [17] A. S. Parkins and H. J. Kimble, quant-ph/9907049.
 - [18] Z. Y. Ou, S. F. Pereira, H. J. Kimble, and K. C. Peng, Phys. Rev. Lett. **68**, 3663 (1992).
 - [19] Z. Y. Ou, S. F. Pereira, and H. J. Kimble, Appl. Phys. B: Photophys. Laser Chem. **55**, 265 (1992).
 - [20] E. P. Wigner, Phys. Rev. **40**, 749 (1932); M. Hillery, R. F. O'Connell, M. O. Scully, and E. P. Wigner, Phys. Rep. **106**, 121 (1984).
 - [21] S. L. Braunstein, A. Mann, and M. Revzen, Phys. Rev. Lett. **68**, 3259 (1992).